The Chat

Vol. 82

Spring 2018

No. 2





The Quarterly Bulletin of the Carolina Bird Club, Inc. The Ornithological Society of the Carolinas

THE CHAT

ISSN No. 0009-1987

Quarterly Bulletin of Carolina Bird Club, Inc.

1909 Lakepark Drive, Raleigh NC 27612

Vol. 82

Spring 2018

No. 2

General Field Notes Editors North Carolina South Carolina

Briefs for the Files

Editor

Christina Harvey William Post

Josh Southern

Steve Shultz 2404 Bristers Spring Way Apex, NC 27523 chat@carolinabirdclub.org

THE CHAT is published quarterly by the Carolina Bird Club, Inc., 9 Quincy Place, Pinehurst NC 28374. Subscription price \$30 per year.

Copyright © 2018 by Carolina Bird Club, Inc. Except for purposes of review, material contained herein may not be reproduced without written permission from the Carolina Bird Club, Inc.

Articles

| 2016 Fall Bird Counts in South Carolina Steven J. Wagner | 48 |
|---|----|
| General Field Notes | |
| First Record of Mountain Plover in North Carolina Jon Altman | 54 |
| Committee Reports | |
| 2017 Annual Report of the North Carolina Bird Records Committee Susan Campbell, et. al | 56 |
| Briefs for the Files | |
| Winter 2017-2018 Josh Southern | |

Cover: Common Eider, 3 January 2018, Dare County, NC Photo by Jeff Beane

2016 Fall Bird Counts in South Carolina

Steven J. Wagner

Department of Liberal Arts Savannah College of Art and Design Savannah, GA 31402 sjwagner@scad.edu

Five Fall Bird Counts were held in 2016. The counts were held September 10 through October 1 in Greenville, Spartanburg, Cherokee, and Aiken counties and in Congaree National Park. The 109 participants recorded a total of 129 species and 10,673 individuals. The species total was well short of the 146 reported last year and the high of 168 in 2011. The 17 species of warblers reported was also well below the 27 reported last year and the high of 29 reported in 2010. Several of the compilers suggested that hot, dry conditions this fall may have contributed their reduced counts.

Greenville County: September 17; 100 species; 3380 individuals

| Of centrine cour | ity: September 17, 100 species, 5500 marriadans |
|------------------|---|
| Compiler: | Paul Serridge |
| | 4 Audrey Lane |
| | Greenville, SC 29615 |
| Participants: | Dave Baker, Joan Baker, Shelby Birch, Andrew Bowers, |
| | Jim Buschur, Marion Clark, Steve Compton, Donnie |
| | Coody, Denise DuPon, Laura Eison, Matthew Eison, |
| | Sherie Gamble, Terry Gamble, Chip Gilbert, Gary |
| | Harbour, Pam Helton, Genny Hill, J. B. Hines, Linda |
| | Hunter, Jerry Johnson, Jo Anne Keasler, David Kirk, Bill |
| | Koon, Len Kopka, Kevin Kubach, Dennis Lee, Tim Lee, |
| | Yves Limpalair, Carole Lyle, Chris Peak, Merikay Pirrone, |
| | Michael Robertson, Barbara Serridge, Paul Serridge, Lily |
| | Shelnut, Tim Shelnut, Larry Sturm, Pat Sturm, Dennis |
| | Trapp, Judith Webb, Anne Williams, Arthur Williams, |
| | Cindy Womble, Louis Womble, Susan Zickos. |
| | |
| Spartanburg Co | ounty: September 24; 84 species; 3954 individuals |
| Compiler: | Lyle Campbell |
| - I | 126 Greengate Lane |
| | Spartanburg, SC 29307-2215 |
| Participants: | Tim Brown, Dan Bryant, Lewanna Caldwell, David |
| Ŧ | Campbell, Lyle Campbell, Sarah Campbell, Timothy |

Campbell, Dan Codispoti, Cheryl Cooksey, Mark

Cooksey, Sharon Donovan, Ed Elam, Kathleen Elam, J. B.

Hines, Frank Hull, Phyllis Hull, Jack Jennings, Jerry Johnson, Don Knight, Mary Knight, Nancy Mandlove, Joe Mitchell, Roberta Mitchell, Nora Moore, Terry Moore, Bill Myers, Mary Ann Myers, Bob Powell, Sandra Powell, Doug Rayner, Mack Shealy, Elizabeth Storm, Jon Storm, Melissa Storm, Kristin Taylor, Gerald Thrumond, M. B. Ulmer, Kay Wealden, Richard Wealden, Dan Wootton, Nancy Wootton

Spartanburg had a fair Fall Count, considering that several key birders were again unable to participate. Birds were scarce, perhaps reflecting weeks of hot, dry weather. We had 84 total species, including only six warbler species. Sparrows and blackbirds were scarce. Four chimneys attracted 1200 migrating Chimney Swifts at dusk. Best birds were Bald Eagle (second year), Gadwall, Cape May Warbler, and Swainson's Thrush.

Cherokee County: September 10; 62 species; 739 individuals

| Compiler: | Lyle Campbell | | | | |
|---------------|--|--|--|--|--|
| | 126 Greengate Lane | | | | |
| | Spartanburg, SC 29307 | | | | |
| Participants: | David Campbell, Lyle Campbell, Sarah Campbell, | | | | |
| | Timothy Campbell, Claude Cobb, Randy Fowler, J. B. | | | | |
| | Hines, Jerry Johnson, Karla Lavender, Janet Tsiknias | | | | |

Ten Cherokee Count birders found 62 species, including only two species of warblers. Birds were scarce, possibly reflecting weeks of hot, dry weather. Our numbers from yard and feeder watches were up. Best bird was a Broad-winged Hawk.

Aiken County: October 1; 83 species; 1528 individuals

| Compiler: | Anne Waters |
|---------------|---|
| | 1621 Apple Valley Dr. |
| | Augusta, GA 30906 |
| Participants: | Meg Francouer, Wade Gassman, Paul Koehler, Mat |
| | Malain, Eric Monaco, Mary Pallon, Lois Stacey, Calvin |
| | Zippler |

Congaree National Park: September 25; 53 species; 1072 individuals

Compiler: John Grego Participants: Ron Ahle, John Grego, Kathleen O'Grady, Alice Steinke, Angela Valasori

A Fall Migration Count was held September 25 at Congaree National Park. Weather was 70 degrees in the morning and 90 degrees in the afternoon, and mosquitoes were active in the floodplain. Thrushes were heard overhead in good numbers from the Visitor Center parking lot predawn, with the nocturnal flight call count dominated by Swainson's Thrush. Birding in the eastern portion of the park was less intensive than usual, which cut down on species diversity in general and migrant counts in particular. 53 species were observed on the day.

| Black-bellied Whistling-Duck 4 4 Canada Goose 173 562 57 32 824 Wood Duck 63 6 32 2 103 Gadwall 2 2 2 3 Mallard 101 115 5 3 224 Am. Black Duck 1 2 3 3 Hoded Merganser 6 36 42 42 Pied-billed Grebe 1 1 4 6 Rock Pigeon 14 123 45 12 194 Eurasian Collared-Dove 1 2 1 4 4 Mourning Dove 187 161 62 41 4 455 Yellow-billed Cuckoo 2 1 1 2 11 1 2 11 1 1 1 12 11 1 1 11 1 12 11 1 1 1 1 12 11 1 1 1 12 11 1 12 11 | | Greenv | Spart | Cherok | Aiken | Congar | Total |
|--|------------------------------|--------|-------|--------|-------|--------|-------|
| Wood Duck 63 6 32 2 103 Gadwall 2 2 Mallard 101 115 5 3 224 Am. Black Duck 1 2 3 3 Hooded Merganser 3 3 3 3 Red-breasted Merganser 6 36 42 Northern Bobwhite 2 1 3 Wild Turkey 6 36 42 Pied-billed Grebe 1 1 4 6 Rock Pigeon 14 123 45 12 194 Eurasian Collared-Dove 1 2 1 4 4 Mourning Dove 187 161 62 41 4 455 Yellow-billed Cuckoo 2 1 1 2 1 </td <td>Black-bellied Whistling-Duck</td> <td></td> <td></td> <td></td> <td>4</td> <td></td> <td>4</td> | Black-bellied Whistling-Duck | | | | 4 | | 4 |
| $\begin{array}{c c c c c c c c c c c c c c c c c c c $ | | | | 57 | | | |
| Mailard 101 115 5 3 224 Am. Black Duck 1 2 3 3 Hooded Merganser 3 6 6 Northern Bobwhite 2 1 3 Wild Turkey 6 36 42 Pied-billed Grebe 1 1 4 6 Rock Pigeon 14 123 45 12 194 Eurasian Collared-Dove 1 2 1 4 45 Mourning Dove 187 161 62 41 4 45 Veilow-billed Cuckoo 2 1 1 5 2 11 Chinney Swift 134 1276 22 14 14 1460 Ruby-throated Hummingbird 722 42 2 3 769 Rufous Hummingbird 1 1 1 1 1 Solitary Sandpiper 2 2 2 4 4 4 Wilson's Phalarope 1 1 1 1 1 1 1 <td></td> <td>63</td> <td></td> <td></td> <td>32</td> <td>2</td> <td></td> | | 63 | | | 32 | 2 | |
| Am. Black Duck 1 2 3 Hooded Merganser 3 3 Red-breasted Merganser 6 3 Northern Bobwhite 2 1 3 Wild Turkey 6 36 42 Pied-billed Grebe 1 1 4 6 Rock Pigeon 14 123 45 12 194 Eurasian Collared-Dove 1 2 1 4 455 Yeld-willed Cuckoo 2 1 1 2 2 11 2 Common Nighthawk 1 1 2 1 1 1 2 14 14 1460 Ruby-throated Hummingbird 722 42 2 3 769 769 Rufous Hummingbird 722 42 2 3 769 769 769 769 769 769 769 78 769 78 769 78 76 72 1 1 1 71 1 71 1 71 1 71 1 71 | | | - | | | | _ |
| Hooded Merganser 3 3 Red-breasted Merganser 6 6 Northern Bobwhite 2 1 3 Wild Turkey 6 36 42 Pied-billed Grebe 1 1 4 6 Rock Pigeon 14 123 45 12 194 Eurasian Collared-Dove 1 2 1 4 455 Yellow-billed Cuckoo 2 1 1 5 2 11 Common Nighthawk 1 1 2 1 1 4 460 Ruby-throated Hummingbird 722 42 2 3 769 79 Rufous Hummingbird 125 21 16 93 1 1 1 Chimey Swift 134 1276 22 14 14 1460 Ruby-throated Hummingbird 722 42 2 3 769 Rufous Hummingbird 1 25 21 16 93 1 1 1 1 1 1 1 1 | | | | 5 | 3 | | |
| Red-breasted Merganser 6 6 6 Northern Bobwhie 2 1 3 Wild Turkey 6 36 42 Pied-billed Grebe 1 1 4 Rock Pigeon 14 123 45 12 194 Eurasian Collared-Dove 1 2 1 4 Mourning Dove 187 161 62 41 4 Mourning Dove 187 161 62 41 4 455 Yellow-billed Cuckoo 2 1 1 5 2 11 Common Nighthawk 1 1 1 2 14 1460 Ruby-throated Hummingbird 722 42 2 3 769 Rufous Hummingbird 1 1 1 1 1 Kildeer 31 25 21 16 93 Least Sandpiper 1 1 1 1 1 Solitary Sandpiper 2 2 2 4 4 Wilson's Phalarope | | 1 | | | | | |
| Northern Bobwhite 2 1 3 Wild Turkey 6 36 42 Pied-billed Grebe 1 1 4 6 Rock Pigeon 14 123 45 12 194 Eurasian Collared-Dove 1 2 1 4 Mourning Dove 187 161 62 41 4 455 Yellow-billed Cuckoo 2 1 1 5 2 11 Common Nighthawk 1 1 2 1 1 2 2 Castern Whip-poor-will 1 1 1 1 1 1 1 1 Common Nighthawk 1 134 1276 22 14 14 1460 Ruby-throated Hummingbird 722 42 2 3 769 Rufous Hummingbird 1 25 21 16 93 Least 3 769 Least Sandpiper 1 1 2 2 4 4 4 Solitary Sandpiper 2 2 <t< td=""><td>e</td><td></td><td></td><td></td><td></td><td></td><td></td></t<> | e | | | | | | |
| Wild Turkey 6 36 42 Pied-billed Grebe 1 1 4 6 Rock Pigeon 14 123 45 12 194 Eurasian Collared-Dove 1 2 1 4 Mourning Dove 187 161 62 41 4 455 Yellow-billed Cuckoo 2 1 1 5 2 11 Common Nighthawk 1 1 2 2 3 769 Ruby-throated Hummingbird 722 42 2 3 769 Rufous Hummingbird 1 1 1 1 1 Killdeer 31 25 21 16 93 2 Least Sandpiper 1 1 2 2 4 4 4 Solitary Sandpiper 2 2 2 4 | e | • | 6 | | | | |
| Picd-billed Grebe 1 1 4 6 Rock Pigeon 14 123 45 12 194 Eurasian Collared-Dove 1 2 1 4 Mourning Dove 187 161 62 41 4 Mourning Dove 187 161 62 41 4 455 Yellow-billed Cuckoo 2 1 1 5 2 11 Common Nighthawk 1 1 1 2 Eastern Whip-por-will 1 1 1 Chimney Swift 134 1276 22 14 14 1460 Ruby-throated Hummingbird 1 1 1 1 1 1 Killdeer 31 25 21 16 93 1 <td></td> <td>2</td> <td></td> <td>-</td> <td></td> <td></td> <td></td> | | 2 | | - | | | |
| Rock Pigeon 14 123 45 12 194 Eurasian Collared-Dove 1 2 1 4 Mourning Dove 187 161 62 41 4 Yellow-billed Cuckoo 2 1 1 5 2 11 Common Nighthawk 1 1 2 1 1 2 Eastern Whip-poor-will 1 1 1 1 1 1 Chinney Swift 134 1276 22 14 14 1460 Ruby-throated Hummingbird 722 42 2 3 769 Rufous Hummingbird 1 1 1 1 1 Killdeer 31 25 21 16 93 Least Sandpiper 1 1 1 1 1 1 Spotted Sandpiper 1 1 1 1 1 1 Solitary Sandpiper 2 2 2 1 1 1 Double-crested Cormorant 1 9 2 1 | 5 | | | 36 | | | |
| Eurasian Collared-Dove1214Mourning Dove18716162414455Yellow-billed Cuckoo2115211Common Nighthawk1112Eastern Whip-poor-will111Chinney Swift13412762214141460Ruby-throated Hummingbird7224223769Rufous Hummingbird1111Killdeer3125211693Least Sandpiper565656Pectoral Sandpiper111Spotted Sandpiper224Lesser Yellowlegs1818Greater Yellowlegs11Duble-crested Cormorant192Anhinga66Great Egret1041327Little Blue Heron298912Black Vulture2698912Green Heron214Cooper's Hawk1214Cooper's Hawk1214Red-shouldered Hawk128565Brack-Shined Hawk128562Red-shouldered Hawk128565Brack Withree2222Black Vulture306725 <td></td> <td>-</td> <td></td> <td></td> <td>-</td> <td></td> <td></td> | | - | | | - | | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | | | 45 | | | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | | | | | | - |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | | | | | | |
| Eastern Whip-poor-will11Chimney Swift13412762214141460Ruby-throated Hummingbird7224223769Rufous Hummingbird1111Killdeer3125211693Least Sandpiper565656Pectoral Sandpiper111Spotted Sandpiper224Lesser Yellowlegs1818Greater Yellowlegs44Wilson's Phalarope11Double-crested Cormorant192Anhinga66Great Egret10413Little Blue Heron223Little Egret7979Green Heron223Black Vulture269891223Bald Eagle112Sharp-shinned Hawk1214214Cooper's Hawk1214222Red-shouldered Hawk55622222Lesser Screech-Owl246Great Horned Owl2482Lesser Yellowlegs33Great Horned Owl246Great Horned Owl246 | | | - | 1 | 5 | 2 | |
| Chinney Swift13412762214141460Ruby-throated Hummingbird7224223769Rufous Hummingbird1111Killdeer3125211693Least Sandpiper5656Pectoral Sandpiper111Spotted Sandpiper224Lesser Yellowlegs1818Greater Yellowlegs11Greater Yellowlegs44Wilson's Phalarope11Double-crested Cormorant19Anhinga66Great Blue Heron8164Great Egret10413Uittle Blue Heron22Black Vulture269891264Turkey Vulture3067250 Soprey123Bald Eagle112Sharp-shinned Hawk1214214Cooper's Hawk1285Broad-winged Hawk2220Eastern Screech-Owl2482482161616 | e | 1 | 1 | | | | |
| Ruby-throated Hummingbird 722 42 2 3 769 Rufous Hummingbird 1 1 1 1 1 Killdeer 31 25 21 16 93 Least Sandpiper 56 56 56 56 Pectoral Sandpiper 1 1 1 1 Spotted Sandpiper 2 2 4 4 Lesser Yellowlegs 18 18 18 Greater Yellowlegs 4 4 4 Wilson's Phalarope 1 1 1 1 Double-crested Cormorant 1 9 2 12 1 Anhinga 6 6 6 6 6 6 Great Blue Heron 8 16 4 5 3 36 Great Egret 10 4 13 27 1 Little Blue Heron 2 2 3 140 Osprey 1 2 3 140 Osprey 1 2 3 3 <t< td=""><td></td><td></td><td></td><td></td><td>-</td><td></td><td>-</td></t<> | | | | | - | | - |
| Rufous Hummingbird 1 1 Killdeer 31 25 21 16 93 Least Sandpiper 56 56 56 56 Pectoral Sandpiper 1 1 1 Spotted Sandpiper 2 2 2 4 Lesser Yellowlegs 18 18 18 Greater Yellowlegs 4 4 Wilson's Phalarope 1 1 1 Double-crested Cormorant 1 9 2 12 Anhinga 6 6 6 Great Blue Heron 8 16 4 5 3 36 Great Egret 10 4 13 27 27 Little Blue Heron 10 10 10 10 Cattle Egret 79 79 79 79 Green Heron 2 1 2 3 140 Osprey 1 2 1 4 Cooper's Hawk 1 2 1 4 Cooper's Hawk 1 | 2 | | | | | 14 | |
| Killdeer3125211693Least Sandpiper5656Pectoral Sandpiper111Spotted Sandpiper224Lesser Yellowlegs1818Greater Yellowlegs44Wilson's Phalarope11Double-crested Cormorant19Anhinga66Great Blue Heron8164Great Egret10413Little Blue Heron22Black Vulture2698Opprey123Blade Eagle11Cosper's Hawk121Arded Hawk222Red-shouldered Hawk121Arded Hawk222Dester Screech-Owl246Great Horned Owl248Starp Screech-Owl24Starp Screech-Owl24 | | | 42 | 2 | 3 | | |
| Least Sandpiper 56 56 Pectoral Sandpiper 1 1 Spotted Sandpiper 2 2 4 Lesser Yellowlegs 18 18 Greater Yellowlegs 4 4 Wilson's Phalarope 1 1 Double-crested Cormorant 1 9 2 12 Anhinga 6 6 6 Great Blue Heron 8 16 4 5 3 36 Great Egret 10 4 13 27 10 10 10 Little Blue Heron 2 79 79 79 79 79 79 Green Heron 2 2 2 3 140 10 10 Cattle Egret 79 79 79 3 3 140 0 0 12 64 14 2 3 3 140 0 0 10 10 10 10 10 10 10 10 10 10 10 11 1 14 10 <td>ę</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> | ę | | | | | | |
| Pectoral Sandpiper111Spotted Sandpiper112Solitary Sandpiper224Lesser Yellowlegs1818Greater Yellowlegs44Wilson's Phalarope11Double-crested Cormorant192Anhinga66Great Blue Heron81645Great Egret1041327Little Blue Heron101010Cattle Egret7979Green Heron226Black Vulture2698912Black Vulture306725153140Osprey123336Great Jagle14233Bald Eagle1214Cooper's Hawk1214Red-shouldered Hawk128565Broad-winged Hawk22202Eastern Screech-Owl2466Great Horned Owl248216 | | 31 | 25 | 21 | | | |
| Spotted Sandpiper112Solitary Sandpiper224Lesser Yellowlegs1818Greater Yellowlegs44Wilson's Phalarope11Double-crested Cormorant192Anhinga66Great Blue Heron81645Great Egret1041327Little Blue Heron101010Cattle Egret7979Green Heron22Black Vulture26989Jadd Eagle112Sharp-shinned Hawk121Cooper's Hawk121Red-shouldered Hawk1285Broad-winged Hawk2220Eastern Screech-Owl248Great Horned Owl248Solar Screech-Owl24Solar Screech-Owl24Sharp-shined Owl24Starp Screech-Owl24Starp | | | | | | | |
| Solitary Sandpiper2224Lesser Yellowlegs1818Greater Yellowlegs44Wilson's Phalarope11Double-crested Cormorant19212Anhinga66Great Blue Heron81645336Great Egret10413271010Little Blue Heron1010101010Cattle Egret79797979Green Heron22264Turkey Vulture306725153140Osprey1233363636Back Vulture1021444Cooper's Hawk12144Red-shouldered Hawk12856536Broad-winged Hawk5562220Eastern Screech-Owl24666Great Horned Owl248216 | | | | | 1 | | |
| Lesser Yellowlegs1818Greater Yellowlegs44Wilson's Phalarope11Double-crested Cormorant192Anhinga66Great Blue Heron81645Great Egret1041327Little Blue Heron101010Cattle Egret7979Green Heron22Black Vulture26989123Bald Eagle112Sharp-shinned Hawk1214Cooper's Hawk1214Red-shouldered Hawk128565Broad-winged Hawk5562220Eastern Screech-Owl248216 | | | 1 | | | | |
| $\begin{array}{cccccccccccccccccccccccccccccccccccc$ | | 2 | | 2 | | | - |
| Wilson's Phalarope11Double-crested Cormorant19212Anhinga66Great Blue Heron816453Great Egret1041327Little Blue Heron1041327Little Blue Heron101010Cattle Egret7979Green Heron22Black Vulture2698912Green Heron23Bladk Vulture306725153140Osprey123Bald Eagle1123Bald Eagle11423Bald Eagle1214Cooper's Hawk1214Red-shouldered Hawk12856536Broad-winged Hawk5562220Eastern Screech-Owl24666Great Horned Owl248216 | | | | | | | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | • | | | | | | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | | | | 1 | | |
| Great Blue Heron81645336Great Egret1041327Little Blue Heron101010Cattle Egret7979Green Heron22Black Vulture2698912Osprey123Bald Eagle112Sharp-shinned Hawk1214Cooper's Hawk1214Red-shouldered Hawk128565Broad-winged Hawk55622Red-tailed Hawk55622Castern Screech-Owl248216 | | 1 | 9 | 2 | | | |
| $\begin{array}{cccccccccccccccccccccccccccccccccccc$ | | | | | | | |
| Little Blue Heron1010Cattle Egret7979Green Heron22Black Vulture269891264Turkey Vulture306725153140Osprey12333Bald Eagle1123Bald Eagle114Cooper's Hawk1214Red-shouldered Hawk12856536Broad-winged Hawk2220220Eastern Screech-Owl2466Great Horned Owl248216 | | | | 4 | | 3 | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | ę | 10 | 4 | | | | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | | | | | | |
| Black Vulture 26 9 8 9 12 64 Turkey Vulture 30 67 25 15 3 140 Osprey 1 2 3 Bald Eagle 1 2 3 Bald Eagle 1 2 1 2 Sharp-shinned Hawk 1 2 1 4 Cooper's Hawk 1 2 1 4 Red-shouldered Hawk 12 8 5 6 5 Broad-winged Hawk 2 2 20 Eastern Screech-Owl 2 4 6 Great Horned Owl 2 4 8 2 | | | | | 79 | | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | | | | | | |
| $\begin{array}{c ccccccccccccccccccccccccccccccccccc$ | | | 9 | | 9 | | |
| Bald Eagle112Sharp-shinned Hawk1214Cooper's Hawk1214Red-shouldered Hawk12856536Broad-winged Hawk222Red-tailed Hawk5562220Eastern Screech-Owl2466Great Horned Owl248216 | Turkey Vulture | | 67 | | 15 | 3 | |
| Sharp-shinned Hawk1214Cooper's Hawk1214Red-shouldered Hawk12856536Broad-winged Hawk222Red-tailed Hawk5562220Eastern Screech-Owl246Great Horned Owl248216 | | 1 | | 2 | | | |
| $\begin{array}{c c} Cooper's Hawk & 1 & 2 & 1 & 4 \\ Red-shouldered Hawk & 12 & 8 & 5 & 6 & 5 & 36 \\ Broad-winged Hawk & 2 & 2 \\ Red-tailed Hawk & 5 & 5 & 6 & 2 & 2 & 20 \\ Eastern Screech-Owl & 2 & 4 & 6 \\ Great Horned Owl & 2 & 4 & 8 & 2 & 16 \\ \end{array}$ | | | | | | 1 | |
| Red-shouldered Hawk 12 8 5 6 5 36 Broad-winged Hawk 2 3 | - | | | | | | |
| Broad-winged Hawk 2 2 2 Red-tailed Hawk 5 5 6 2 2 20 Eastern Screech-Owl 2 4 6 6 Great Horned Owl 2 4 8 2 16 | 1 | | | | | | 4 |
| Red-tailed Hawk 5 5 6 2 2 20 Eastern Screech-Owl 2 4 6 6 Great Horned Owl 2 4 8 2 16 | | 12 | 8 | | 6 | 5 | 36 |
| Eastern Screech-Owl246Great Horned Owl248216 | Broad-winged Hawk | | | 2 | | | 2 |
| Great Horned Owl 2 4 8 2 16 | | | 5 | 6 | | 2 | |
| | | | | | | | 6 |
| Barred Owl 3 1 1 9 15 29 | | | | | | | |
| | Barred Owl | 3 | 1 | 1 | 9 | 15 | 29 |

Table 1. Fall 2016 Bird Counts in South Carolina

50

| | Creamy | Crocet | Chanalr | Ailton | Concor | Total |
|--------------------------|-------------|-------------|-------------|------------|--------|-------------|
| Belted Kingfisher | Greenv 7 | Spart 13 | Cherok 3 | Aiken 4 | Congar | Total 27 |
| Red-headed Woodpecker | 7 | 3 | 2 | 16 | 16 | 44 |
| Red-bellied Woodpecker | 41 | 33 | 13 | 34 | 23 | 144 |
| Downy Woodpecker | 29 | 30 | 2 | 18 | 10 | 89 |
| Hairy Woodpecker | 5 | 1 | 1 | | 1 | 8 |
| Red-cockaded Woodpecker | U | • | - | 2 | • | 2 |
| Northern Flicker | 3 | 5 | 3 | 13 | 3 | 27 |
| Pileated Woodpecker | 33 | 6 | 4 | 23 | 15 | 81 |
| American Kestrel | 2 | 2 | 1 | 4 | | 9 |
| Peregrine Falcon | 1 | _ | - | 1 | | 2 |
| Eastern Wood-Pewee | 15 | 6 | 5 | 8 | 2 | 36 |
| Acadian Flycatcher | 1 | 2 | | | 3 | 6 |
| Eastern Phoebe | 29 | 15 | 9 | 11 | 1 | 65 |
| Great Crested Flycatcher | | | 2 | | | 2 |
| Eastern Kingbird | 1 | 1 | _ | | | 2 |
| White-eyed Vireo | 32 | 4 | 8 | 29 | 16 | 89 |
| Yellow-throated Vireo | 2 | 1 | | | 2 | 5 |
| Blue-headed Vireo | 5 | | | | | 5 |
| Red-eyed Vireo | 11 | 2 | | | 1 | 14 |
| Blue Jay | 116 | 81 | 23 | 77 | 9 | 306 |
| American Crow | 306 | 149 | 47 | 57 | 12 | 571 |
| Fish Crow | 16 | 2 | 3 | 5 | | 26 |
| crow sp. | 10 | - | U | U | 2 | 2 |
| Common Raven | 5 | | | | _ | 5 |
| Tree Swallow | 6 | | | 9 | | 15 |
| N. Rough-winged Swallow | 0 | | | 165 | 2 | 167 |
| Barn Swallow | | 1 | | 16 | - | 17 |
| Carolina Chickadee | 100 | 98 | 32 | 41 | 12 | 283 |
| Tufted Titmouse | 78 | 85 | 35 | 75 | 30 | 303 |
| White-breasted Nuthatch | 39 | 19 | 7 | 16 | 10 | 91 |
| Brown-headed Nuthatch | 10 | 12 | 1 | 15 | 6 | 44 |
| House Wren | 2 | 12 | | 2 | | 16 |
| Carolina Wren | 91 | 60 | 26 | 59 | 45 | 278 |
| Blue-gray Gnatcatcher | 6 | | | | 1 | 7 |
| Eastern Bluebird | 119 | 47 | 20 | 10 | | 196 |
| Veery | | | | | 88 | 88 |
| Gray-cheeked Thrush | | | | | 5 | 5 |
| Swainson's Thrush | 1 | 3 | | | 571 | 575 |
| Wood Thrush | | | | | 19 | 19 |
| American Robin | 74 | 53 | 23 | | 1 | 151 |
| Gray Catbird | 16 | 3 | | 3 | | 22 |
| Brown Thrasher | 26 | 21 | 1 | 22 | 1 | 71 |
| Northern Mockingbird | 50 | 52 | 8 | 32 | | 142 |
| European Starling | 52 | 53 | 58 | 115 | | 278 |
| Cedar Waxwing | 15 | | | | | 15 |
| House Sparrow | 2 | 20 | 2 | | | 24 |
| House Finch | 55 | 70 | 10 | 2 | | 137 |
| Purple Finch | | 2 | | | | 2 |
| American Goldfinch | 133 | 30 | 8 | 2 | 1 | 174 |
| Eastern Towhee | 17 | 5 | 6 | 21 | | 49 |
| Chipping Sparrow | 4 | 14 | 10 | 2 | | 30 |
| Field Sparrow | 2 | 14 | 8 | | | 24 |
| Song Sparrow | 3 | 6 | | | | 9 |
| Dark-eyed Junco | 12 | | | | | 12 |
| Eastern Meadowlark | 1 | | 2 | | | 3 |
| Baltimore Oriole | 1 | | | | | 1 |
| Red-winged Blackbird | | 1 | 1 | 1 | | 3 |
| 0 | | | | | | |

2016 Fall Bird Counts in South Carolina

| Brown-headed Cowbird | Greenv | Spart 133 | Cherok | Aiken 2 | Congar | Total 138 |
|---|-------------|--------------|-----------|------------|------------|--------------|
| Common Grackle | 18 | 129 | 8 | 4 | | 159 |
| Ovenbird | 1 | 1 | | | 1 | 1 |
| Worm-eating Warbler Northern Waterthrush | | 1 | | | 1 | 2 |
| Black-and-white Warbler | 2 | | | | 1 2 | 1 4 |
| Tennessee Warbler | 2 9 | | | | Z | 4 9 |
| Common Yellowthroat | 8 | | | 4 | 5 | 17 |
| Hooded Warbler | ° 5 | | | 4 | 2 | 8 |
| American Redstart | 22 | 3 | 2 | 4 | 18 | 8 49 |
| Cape May Warbler | 1 | 1 | 2 | 4 | 18 | 49 |
| Northern Parula | 7 | 1 | | 11 | 19 | 37 |
| Magnolia Warbler | 3 | | | 4 | 19 | 8 |
| Yellow Warbler | 1 | 1 | | 4 | 1 | 2 |
| Chestnut-sided Warbler | 6 | 1 | | 2 | | 2 8 |
| Black-throated Blue Warbler | 1 | | | 2 | | 0 1 |
| Palm Warbler | 15 | | | 12 | | 27 |
| Pine Warbler | 9 | 11 | 8 | 49 | 16 | 93 |
| Yellow-throated Warbler | 2 | 1 | 0 | 2 | 10 | 93 6 |
| Summer Tanager | 4 | 4 | 2 | 3 | 6 | 19 |
| Scarlet Tanager | 8 | 1 | 1 | 5 | 1 | 11 |
| Northern Cardinal | 105 | 160 | 17 | 83 | 15 | 380 |
| Rose-breasted Grosbeak | 105 | 2 | 17 | 65 | 15 | 3 |
| Blue Grosbeak | 3 | 2 | 2 | 3 | | 8 |
| Indigo Bunting | 26 | 7 | 3 | 15 | 1 | 52 |
| Painted Bunting | 20 | , | 5 | 1 | 1 | 1 |
| T anneed Dunting | | | | 1 | | 1 |
| Species Individuals | 100 3380 | 84 3954 | 62 739 | 83 1528 | 53 1072 | 129 10673 |
| liferviduals | 5580 | 3934 | 139 | 1528 | 1072 | 10075 |
| Describer Count | | | | | | |
| Regular Count Hours foot | | 30 | 10 | 14.3 | 10 | 64.3 |
| Hours car | | 12 | 8.5 | 14.5 | 10 | 38.35 |
| Hours canoe/kayak | | 12 | 0.5 | 3 | | 38.33 |
| Hours boat | | 7.5 | | 5 | | 7.5 |
| Hours bicycle | | 0.5 | | 3 | | 3.5 |
| Hours bleyele | | 0.5 | | 5 | | 5.5 |
| Miles foot | | 17.5 | 11.5 | 4.5 | 4.5 | 38 |
| Miles car | | 343 | 174 | 111.45 | | 628.45 |
| Miles canoe/kayak | | 0.0 | 171 | 2 | | 2 |
| Miles boat | | 17 | | - | | 17 |
| Miles bicycle | | 6 | | 4 | | 10 |
| | | | | - | | |
| # Regular parties | 16 | 15 | 7 | 6 | | 44 |
| # Regular observers | 45 | 21 | 8 | 8 | 5 | 87 |
| # Species | | 80 | 61 | 79 | | |
| # Individuals | | 2184 | 653 | 1482 | | |
| | | | | | | |
| Stationary | | | | | | |
| Hours Stationary | | | | 2 | | 2 |
| # Parties Stationary | | | | 1 | | 1 |
| # Observers Stationary | | | | 1 | | 1 |
| # Species | | | | 15 | | 15 |
| # Individuals | | | | 24 | | 24 |

| | 1 0 | | | | | |
|-------------------------|--------|--------|----------|----------|--------|--------|
| | Greenv | Spart | Cherok | Aiken | Congar | Total |
| Feeder Watch | | | | | | |
| Hours Feeder Watch | | 52 | 8.5 | | | 60.5 |
| # Feeder Watchers | | 28 | 3 | | | 31 |
| # Feeder Stations | | 17 | 3 | | | 20 |
| # Species | | 39 | 26 | | | |
| # Individuals | | 564 | 85 | | | 649 |
| Nocturnal | | | | | | |
| Hours Nocturnal | | 4 | 0.5 | 4.25 | 1.25 | 10 |
| Miles Nocturnal | | 12 | 0 | 8.5 | 5 | 25.5 |
| # Parties Nocturnal | | 4 | 1 | 3 | | 8 |
| # Observers Nocturnal | | 6 | 1 | 3 | | 10 |
| # Species | | 4 | 1 | 7 | | |
| # Individuals | | 1206 | 1 | 30 | | 1237 |
| Total | | | | | | |
| Party Hours | 75.5 | 102 | 27.5 | 44.4 | 11.25 | 260.65 |
| Party Miles | | 383.5 | 185.5 | 130.45 | 9.5 | 708.95 |
| T | 0.000 | 0700 | 0700 | 0515 | 0520 | |
| Time start | 0600 | 0700 | 0700 | 0515 | 0530 | |
| Time stop | 1930 | 2200 | 2100 | 1900 | 1825 | |
| Temperature: pre-dawn | | 70 | 68 | 62 | | |
| dawn | | 69 | 68 | 62 | 70 | |
| AM | 61 | 77 | 80 | 69 | | |
| noon | | 81 | 88 | 76 | | |
| PM | 86 | 87 | 95 | 86 | 90 | |
| sunset | | 80 | 80 79 | 80 79 | | |
| night | | 77 | 78 | 78 | | |
| wind: pre-dawn | | | | 0 | | |
| dawn | <5 | | | 0 | | |
| AM | <5 | | | SW3 | | |
| noon | <5 | | | SW5 | | |
| PM | <5 | | | WSW7 | | |
| sunset | <5 | | | W5 | | |
| night | | | | W5 | | |
| %clouds: pre-dawn | | 0 | 0 | 20 | | |
| dawn | | 0 | 0 | 20 | | |
| AM | | 0 | 0 | 20 | | |
| noon | | 0 | 0 | 10 | | |
| PM | | 0 | 0 | 15 | | |
| sunset | | 0 | 0 | 20 | | |
| night | | 0 | 0 | 20 | | |
| precipitation: pre-dawn | 0 | 0 | 0 | 0 | | |
| dawn | 0 | 0 | 0 | 0 | | |
| AM | 0 | 0 | 0 | 0 | | |
| noon | 0 | 0 | 0 | 0 | | |
| PM | 0 | 0 | 0 | 0 | | |
| sunset | 0 | 0 0 | 0 0 | 0 0 | | |
| night | | 0 | U | U | | |

The Chat, Vol. 82, No. 2, Spring 2018

Jon Altman

131 Charles Street Harkers Island, NC 28531 jon_altman@nps.gov

On January 26, 2016, I found an unusual plover with a buffy tinge to the breast at Cape Lookout National Seashore, Carteret County, NC. The plover, noticeably larger than the expected *Charadrius* species, was roosting on a large sandflat on the southern shore of Ophelia Inlet on South Core Banks. The purpose of my visit to this site was to survey for Piping Plover (*Charadrius melodus*), Wilson's Plover (*Charadrius wilsonia*), Snowy Plover (*Charadrius alexandrinus*) and Red Knot (*Calidris cantus*) as part of the 2016 International Piping Plover Census.

After ruling out the survey species, consulting the Ibird application, and studying the bird from a distance of 20 meters with a spotting scope, I believed that the unusual bird was a Mountain Plover (*Charadrius montanus*). The tide was high during the observation period, and all shorebirds were roosting. The Mountain Plover made some minor movements, but primarily rested near some debris with other shorebirds. Piping and Wilson's plovers are common birds of the seashore, with 16 Piping Plovers recorded at this wintering site during the census. Snowy Plovers are infrequent summer visitors. Though similar in overall color and common as a wintering species, Semipalmated Plover (*Charadrius semipalmatus*) was eliminated due to its orange legs and smaller size. Photographs were taken for further study and documentation (Figure 1).

Back in the office, I consulted The Sibley Guide to Birds (Sibley 2000) to confirm species identification and range. Close inspection of the pictures appeared to confirm my initial identification of Mountain Plover, and lack of orange/buffy fringes to the feathers of the mantle and scapulars suggested that the bird was a non-breeding adult. Gender could not be determined. Information on the sighting and photographs of the bird were submitted to The Carolina Bird Club website's photo gallery. Multiple observers ventured out to Ophelia Inlet to relocate the bird and pictures were shared of re-sightings on January 29 and 31, 2016 and February 1 and 9, 2016.

The Mountain Plover is a bird of interior and western North America. It breeds in open, flat, disturbed areas from Alberta and Saskatchewan to Texas and winters primarily in California's Central Valley. It is a described as a vagrant along the East Coast (Paulson, 2005). Haymen et. al., (1986) reports sightings in Florida, Virginia, and Massachusetts. More recently a sighting from October 10, 2007 at Huntington Beach State Park, SC was accepted by the South Carolina Bird Records Committee (Slyce, et al, 2009). There is no prior record of this species in North Carolina.

Literature Cited:

Haymen, P., J. Marchant, and T. Prater. 1986. Shorebirds: An Identification Guide. Houghton Mifflin, Massachusetts.

Paulson, Donald. 2005. Shorebirds of North America: The Photographic Guide. Princeton University Press, New Jersey.

Sibley, D.A. 2000. National Audubon Society: The Sibley Guide to Birds. Alfred A. Knopf, New York.

Slyce D., G. Beaton, L. Glover, C. Hill, T. Piephoff, W. Post, and S. Wagner. 2009. 2008 Annual Report of the South Carolina Bird Records

Committee. Chat 73:105-106.



Figure 1. Mountain Plover (*Charadrius montanus*) roosting with a Semipalmated Plover (*Charadrius semipalmatus*) and Least Sandpiper (*Calidris minutilla*) at Ophelia Inlet, NC on January 26, 2016. Photo by Jon Altman.

2017 Annual Report of the North Carolina Bird Records Committee

Susan M. Campbell¹, Chair, Keith E. Camburn, Samuel Cooper, W. Jefferson Lemons, Michael W. D. McCloy, Jeffrey S. Pippen, Simon R. B. Thompson, Joshua L. Southern, Robert W. Van Epps

> ¹ P.O. Box 1443, Southern Pines, NC 28388 susan@ncaves.com

The make-up of the 2017 North Carolina Bird Records Committee included additions of Keith Camburn, Jeffrey Pippen, and Robert Van Epps as voting members; and the retention of Richard J. Davis and Harry E. LeGrand Jr. in advisory capacities. Taylor Piephoff, Steven Shultz, and Nathan Swick rotated off the Committee for at least a one-year hiatus.

Accepted Reports

Anna's Hummingbird (*Calypte anna*) (17-01). A written description and photos of an adult male banded in Buxton (Dare) on 22 January 2017 were submitted by Susan Campbell. The bird was first observed by Ann Maddock on 24 January 2017 and persisted through 15 March. Both the written material as well as the photos were accepted unanimously by the committee. This represents the fourth record for the state.

Trumpeter Swan (*Cygnus buccinator*) (17-02). Two written descriptions (Simon Thompson and John Koon) as well as photos were submitted of a bird on the French Broad River (Buncombe) in mid-February 2017. The committee voted unanimously to accept all of the material. This record is the eighth for NC and the first for the mountains.

Arctic Tern (*Sterna paradisaea*) (17-03). Written details and photos were submitted by Ed Corey for a bird present at Falls Lake (Wake) on 25-26 May 2017. The committee voted unanimously to accept the material. This record constitutes the third Piedmont record and the first from that region to be documented by photos.

Tropical Kingbird (*Tyrannus melancholicus*) (17-04). John Fussell submitted both a written report as well as a photo of a bird located at North River Farm (Carteret) on 3 June 2017. This record was approved (8-1). It represents a first for the summer season in North Carolina.

Swinhoe's Storm-Petrel (*Oceanodroma monorhis*) (17-05). Brian Patteson located an individual 30 nm SE of Hatteras (Dare) on 9 June 2017.

Both written details and a photo were submitted. The committee voted unanimously to accept this report, the fourth state record.

Long-billed Dowitcher (*Limnodromus scolopaceus*) (17-06). Wayne Forsythe submitted a written report as well as photos for a bird found on 22 July 2017 at Hendersonville (Henderson). Though twice as many members accepted the photos as did not (6-3), such a voting verdict does not accept a record; there must be no more than one non-accept vote for approval. Fortunately, the written material was approved unanimously, thus providing the first mountain state record for the species.

Sooty Tern (*Onychoprion fuscatus*) (17-08). Simon Thompson located an individual at Hooper Lane (Henderson) on 13 September 2017. A written report and photos were submitted to the committee. The material was accepted unanimously. This record is the first for the mountain region of the state.

Bell's Vireo (*Vireo bellii*) (17-10). James Poling submitted both written details as well as photos for a bird at Warren Wilson College (Buncombe) from 14 November until at least 16 November 2017. The material was accepted by the committee (unanimously for the photos but 6-2 for the written report). This record represents the seventh for the state and the first for the NC mountains.

Black-legged Kittiwake (*Rissa tridactyla*) (17-11). Dwayne Martin submitted both a written report and photo for an individual found on Lake Hickory (Catawba) on 17 November 2017. The committee voted unanimously to accept the record. This record is the second for the species in the Piedmont region and is the third inland record in the state.

Broad-billed Hummingbird (*Cynanthus latirostris*) (17-12). A written description and photos of an adult female banded in Southern Shores (Dare) on 19 November 2017 were submitted by Susan Campbell. The bird was first observed by Jeff Lewis on 30 October 2017 and was present until 1 January 2018, when it was found deceased under the feeder, apparently unable to survive the unusual cold spell that extended all the way to the coast that week. The committee voted unanimously to accept both the written report and the photos. This record is the fourth for the state and is the first female of the species documented here.

Brown Booby (*Sula leucogaster*) (17-13). A written report was received from Ricky Davis of a bird seen on 18 December 2016 at Fort Macon State Park (Carteret). The committee voted unanimously to accept the report. This represents the first winter record for NC (Southern 2017).

Non-Accepted Reports

American Flamingo (*Phoenicopterus ruber*) (17-09). A vote was taken on a written description of five individuals seen flying offshore from a cruise ship off Core Banks (Carteret) on 6 November 2017. Unfortunately no photos were obtained and the description was vague. The committee voted

6-1 not to accept the report, given the insufficient details, especially for what would have been a first record for North Carolina.

Unresolved

There are no unresolved reports as the result of voting during 2017. However, two reports - Yellow-legged Gull (*Larus michahellis*) and Common Swift (*Apus apus*) - were received late in the year that are expected be taken up by the committee in 2018; neither of these species is on the state list.

A list of all the species reviewed can be seen at: http://www.carolinabirdclub.org/brc/NC/reports.html.

Summary

This year's voting resulted in a number of seasonal firsts, as well as four first records for the mountains of North Carolina. Even though no changes in status of Accepted species occurred as a result of committee activity, the list has been reduced by one. The decision by the American Ornithological Society to lump Thayer's Gull with Iceland Gull (Fifty-eighth supplement to the American Ornithological Society's Check-list of North American Birds) now means there are no longer 482 but 481 species on the state list (466 Definitive, 13 Provisional, and two Not Established).

Literature Cited

Chesser, R. Terry, Kevin J. Burns, Carla Cicero, Jon L. Dunn, Andrew W. Kratter, Irby J. Lovette, Pamela C. Rasmussen, J. V. Remsen, Jr., James D. Rising, Douglas F. Stotz, and Kevin Winker. 2017. Fifty-eighth supplement to the American Ornithological Society's Check-list of North American Birds. Auk 134(3):751-773.

LeGrand, H., J. Haire, N. Swick, and T. Howard. 2017. Birds of North Carolina: their Distribution and Abundance.

Southern, J. 2017. Briefs for the Files. Chat 81:49.



H Douglas Pratt

BRIEFS FOR THE FILES

Josh Southern

203 Hyannis Drive Holly Springs, NC 27540 joshsouthern79@gmail.com

(All dates Winter 2017-2018, unless otherwise noted)

Briefs for the Files is a seasonal collection of uncommon-to-rare or unusual North and South Carolina bird sightings and events which do not necessarily require a more detailed Field Note or article. Reports of your sightings are due the 20th of the month after the end of the previous season.

| Winter | December 1–February 28 | due March 20 |
|--------|------------------------|-----------------|
| Spring | March 1–May 31 | due June 20 |
| Summer | June 1–July 31 | due August 20 |
| Fall | August 1–November 30 | due December 20 |

Reports may be submitted in any format, but I prefer that you use email, list multiple sightings in taxonomic order (rather than by date or location), and type your report directly into the body of the email. If your sightings are in a file, please copy-and-paste the text into the body of the email, rather than sending an attachment.

Suitable reports for the Briefs include any sightings you feel are unusual, rare, noteworthy, or just plain interesting to you in any way! It is my responsibility to decide which reports merit inclusion in the Briefs.

Please be sure to include details of any rare or hard-to-identify birds.

I rely in part on sightings reported in Carolinabirds. Please don't, however, rely on me to pick up your sightings from Carolinabirds. Instead, please also send your sightings directly to me as described above.

If I feel that your sighting warrants a Field Note, I will contact either you or the appropriate state Field Notes editor. You may, of course, submit your Field Note directly to the editor without going through me.

Abbreviations: **BRC** – Bird Records Committee, **CBC** – Christmas Bird Count, **Co** – County, **et al.** – and others, **Ft** – Fort, **m. obs.** – multiple observers, **Mt** – Mount, **NC** – North Carolina, **NF** – National Forest, **NWR** – National Wildlife Refuge, **Rd** – Road, **SC** – South Carolina, **SP** – State Park, **St** – Street, **WMA** – Wildlife Management Area, **WTP** – Water Treatment Plant **Snow Goose**: Several small flocks were seen in the southern NC mountains this winter, including up to nine at Carolina Lake, Transylvania Co, 21 Dec through 1 Jan (Kevin Lash, m. obs.) and six at Mills River WTP, Henderson Co, NC, 17 Feb (Denise Hargrove). Individuals in Wilkes Co, 3 Dec (Dwayne Martin) through 5 Jan (Guy McGrane), at the Biltmore Estate, Buncombe Co, 11 Jan through 17 Feb (Doug Johnston, m. obs.); and at Lake Junaluska, Haywood Co, 2 Feb (Bob Olthoff, Connie & Stan Wulkowicz, et al.) were also unusual for the NC mountains.

Ross's Goose: Atypically, in NC, more sightings were made in the Mountain region than in the Coastal Plain this winter. One seen at the Biltmore Estate, Buncombe Co, throughout December, was joined by three other small white geese, either Ross's Geese or Snow x Ross's Goose hybrids, 11 Jan (Doug Johnston, m. obs.). Two were seen on Lake Junaluska, Haywood Co, 2-17 Jan (Bob Olthoff, Connie & Stan Wulkowicz, m. obs.). Two were seen in agricultural fields along US-64 in Tyrell Co and Washington Co, 15 Jan (Ed Corey) through 3 Feb (Ricky Davis). One photographed at the Cedar Island Ferry Terminal, Carteret Co, 5 Dec (Chandra Biggerstaff) was unusual for that area.

Greater White-fronted Goose: High counts included 14 at Pettigrew SP, Washington Co, NC, during its CBC, 27 Dec (Al Hooks, Guy McGrane); 12 at Mattamuskeet NWR, Hyde Co, NC, during its CBC, 29 Dec (Steve Shultz); 17 in Polk Co, NC, where photographed, 13 Jan (Michael Robertson); 15 at Lake Leroy, Oconee Co, SC, 21 Jan (Mark Hopey); 11 in the fields where US-64 crosses over Beasley Rd in Washington Co, NC, 17 Feb (Jeff Lewis); and 27 at Dobbins Farm in Anderson Co, SC, 23 Feb (Jo Anne Keasler). Two hung around Kale Lake in Spartanburg Co, SC, 30 Dec (David Campbell) into March (m. obs.).

Cackling Goose: Three at the Beaufort Club golf course in Carteret Co, NC, 16 Dec (Martin Wall) were a first for that county. Other sightings included one in Forsyth Co, NC, 15 Dec through 2 Jan (m. obs.); one along the Nags Head causeway, Dare Co, NC, 1-4 Jan (Jeff Lewis, m. obs.); two in Guilford Co, NC, 3-7 Jan (Henry Link, m. obs.); and one to five where US-64 crosses Beasley Rd in Washington Co, NC, 9 Jan through 23 Feb (m. obs.).



Cackling Geese (with Canada Goose) 16 December 2017 Carteret Co, NC Photo by Martin Wall **Tundra Swan**: Six on Salem Lake, Forsyth Co, NC, 26 Dec (David & Susan Disher, m. obs.) were unusual for that area.

Eurasian Wigeon: Mattamuskeet NWR and Pea Island NWR, two of the more reliable sites to find this species in NC, hosted, respectively, one to three 14 Dec (Phil Doerr) through 18 Feb (Jamie Adams), and one to two 2 Dec (Al Hooks, Jeff Lewis) through early February (m. obs.). In SC, individuals were seen at Bear Island WMA, Colleton Co, 23 Dec (Ed Blitch, Pam Ford, Jim Pope) through 19 Feb (Steve Compton) and at Santee Coastal Reserve, Charleston Co, 8 Feb (Ritch Lilly).

American Wigeon: 30,683 in Hyde Co, NC, during the Mattamuskeet NWR CBC, 29 Dec (*fide* Susan Campbell) was the highest total on any 2017 Audubon CBC and a record total for that count.

Northern Pintail: 113,521 in Hyde Co, NC, during the Mattamuskeet NWR CBC, 29 Dec (*fide* Susan Campbell) was the highest total on any 2017 Audubon CBC and a record total for that count. 50 in Transylvania Co, NC, 15 Jan (Wayne Forsythe) was a notable count for the mountain region.

"Common Teal": A Eurasian race Green-winged Teal was seen at Mattamuskeet NWR, Hyde Co, NC, 3 Feb (Clyde Sorenson, et al.) and 8 Feb (Jeff Lewis).

Tufted Duck: Pending review by the SC BRC is the report of an apparent Tufted Duck seen and photographed on Bulls Island, Cape Romain NWR, Charleston Co, SC, 2 Feb (David McLean, David Youngblood) into spring (m. obs.). If accepted, this report would provide the first record of this Eurasian species in SC.

Lesser Scaup: 9000 on Stumpy Point Bay, Dare Co, NC, during the Alligator River NWR CBC, 30 Dec (Jay Ross, *fide* Jeff Lewis) was a noteworthy count.

King Eider: Pending review by the SC BRC is the report of an apparent hen King Eider seen and photographed at the 2^{nd} Avenue Pier in Myrtle Beach, Horry Co, SC, 7 Jan (Ritch Lilly) through 6 Feb (m. obs.).

Common Eider: Sightings included one at the pier at Myrtle Beach SP, Horry Co, SC, 19 Nov through 28 Feb (William Kidwell, m. obs.); an adult drake at the Folly Beach Fishing Pier, Charleston Co, SC, 23 Nov through 18 Dec (Simon Harvey, m. obs.); two at Johnnie Mercer's Pier, New Hanover Co, NC, 1-3 Dec (Jamie Adams, m. obs.); two at Jennette's Pier, Dare Co, NC, 16 Dec (Ricky Davis); two at Oregon Inlet, 1-3 Jan (Jeff Lewis, m. obs.); and three in Beaufort Inlet off Ft Macon SP, Carteret Co, NC, 18 Feb (Martin Wall).

Harlequin Duck: An adult drake seen and photographed at Jennette's Pier, Dare Co, NC, 30-31 Dec (m. obs.) was probably the same adult drake seen around the south end of the nearby Bonner Bridge at Oregon Inlet, 1-3 Jan (m. obs.). It's also possible that the adult drake seen and photographed around the jetty at Murrells Inlet, Georgetown Co, SC, 7-8 Jan (Ritch Lilly, m. obs.) was that same individual. Elsewhere, two hens were seen off the Coquina rock outcropping at Ft Fisher, New Hanover Co,

NC, 29 Dec (Ron Clark, Matt Janson) and a first-year drake was photographed at the ferry dock in Hatteras, Dare Co, NC, 3 Jan (Jeff Beane, Stephanie Horton).

Surf Scoter: Ten on Salem Lake, Forsyth Co, NC, 15 Dec (John Haire, Marbry Hopkins) was a good count for an inland site. Other inland sightings included one on Lake Junaluska, Haywood Co, NC, 2 Dec (Connie & Stan Wulkowicz, et al.); two on Lake Townsend, Guilford Co, NC, 11 Dec (Henry Link, m. obs.); and one in Barnwell Co, SC, during the Savannah River Site CBC, 14 Dec (Peter Stangel).

White-winged Scoter: Inland sightings included 12 on Lookout Shoals Lake, Catawba Co, NC, 6 Jan (Dwayne Martin); up to eight on Oak Hollow Lake, Guilford Co, NC, 7 Jan through 15 Feb (Roger Shaw, m. obs.); three on Salem Lake, Forsyth Co, NC, 7 Feb (John Haire); two on Jordan Lake, Chatham Co, NC, 9 Feb (Matt Spangler); and two near the dam on Lake Gaston, Northampton Co, NC, 11 Feb (Ricky Davis).

Long-tailed Duck: There was an influx of this species on inland lakes in January, likely due to below-freezing temperatures at sites to the north. A few of the higher counts were seven on Lake Rhodhiss, Burke Co, 13 Jan (Michael Robertson); six on Oak Hollow Lake, Guilford Co, 14-16 Jan (Mike Turner, m. obs.); 22 on Salem Lake, Forsyth Co, 14 Jan (John Haire); and five on Jordan Lake, Chatham Co, 26 Jan (Matt Spangler). Sightings made at inland sites in SC included one at Dobbins Farm, Anderson Co, 10 Dec (George McHenry); two on Lake Wateree, Kershaw Co, 28 Dec through 5 Jan (Steve Patterson, m. obs.); three at Arcadia Farms in Orangeburg Co, 15 Jan (Julie Mobley); three on Lake Murray, Lexington Co, 6 Feb (Jennifer Shockley); and two on a pond in Ridgeville, Dorchester Co, 17-27 Feb (Tracee Clapper, m. obs.). Four around the Charleston Battery, Charleston Co, 5-20 Jan (Craig Watson, m. obs.) was a good number for a site so far south.

Common Goldeneye: This duck was well-reported after a period of below-normal temperatures in early January. High counts included five on a pond along SC-546 in Greenville Co, SC, 4-6 Jan (Yves Limpalair, m. obs.); 11 on Lake Hickory in Caldwell Co, NC, 10 Jan (Dwayne Martin); five on Lake Townsend, Guilford Co, NC, 10-13 Jan (Matt Wangerin, m. obs.); eight on Beaverdam Reservoir, Wake Co, NC, 21 Jan (Harry LeGrand); 25 on the Cape Fear Fisher off the Ft Fisher Air Force Recreation Area, 3 Feb (Sam Cooper); ten on the Roanoke River in Northampton Co, NC, 11 Feb (Ricky Davis); and six on a pond in Wilkesboro, Wilkes Co, NC, 18 Feb (Guy McGrane).

Hooded Merganser: 1200 crowded into an unfrozen portion of Lake Crabtree, Wake Co, NC, following a period of below-freezing temperatures, 7 Jan (Harry LeGrand) was a notable concentration.

Common Merganser: Like several other species of diving ducks, Common Mergansers were well-reported on lakes and ponds following a period of below freezing temperatures in early January. A few of the higher counts, all from NC, were 15 on Lake Brandt, Guilford Co, 4 Jan (Matt Wangerin); six on Lake Junaluska, Haywood Co, 4 Jan (Vince Kloster); 15 on W. Kerr Scott Reservoir, Wilkes Co, 5 Jan (Guy McGrane, et al.); seven on Lake Crabtree, Wake Co, 8-11 Jan (Eddie Owens, m. obs.); seven on Beaverdam Reservoir, Wake Co, 11 Jan (Harry LeGrand); 15 at Swann WTP, Forsyth Co, 10-18 Feb (Shelley Rutkin, m. obs.); and six on Lake Gaston, Northampton Co, 11 Feb (Ricky Davis). A hen on Crystal Lake in Horry Co, SC, 5-29 Jan (Frank Lawkins, m. obs.) was unusual for a site so far south.

Red-necked Grebe: This species was well-reported this winter. Three photographed together near the southern end of the Bonner Bridge, Dare Co, NC, 15 Jan (Joe Donahue, et al.) and two off Coquina Beach, Dare Co, NC, 21 Jan (Jeff Lewis) were notable, as most sightings made in the Carolinas are of individuals. Individuals were seen on Lake Townsend, Guilford Co, NC, 17 Nov through 5 Jan (Henry Link, m. obs.); at Cape Point, Dare Co, NC, 15 Dec through 17 Feb (m. obs.); off Mitchellville Beach Park, Beaufort Co, SC, 25 Dec (Andy Martin); on Lake Hickory, border of Alexander Co and Catawba Co, NC, 26-27 Dec (Dwayne Martin); at Oregon Inlet, Dare Co, NC, 14 Jan through 16 Feb (m. obs.); off Jennette's Pier, Dare Co, NC, 20-24 Jan (m. obs.); at Murrells Inlet, Georgetown Co, SC, 25 Jan (Chris Clack); on the Cape Fear River near Ft Fisher, New Hanover Co, NC, 3 Feb (Sam Cooper); on Jordan Lake, Chatham Co, NC, 26-27 Feb (Martin Wall, et al.).

Eared Grebe: Individuals were seen at Masonboro Inlet, New Hanover Co, NC, 30 Dec (Sam Cooper, et al.); at Cape Point, Dare Co, NC, 9 Jan (Brian Patteson), 17 Feb (Andrew Thornton) and 23 Feb (Jack Rogers); on the Cape Fear River near Ft Fisher, New Hanover Co, NC, 21 Jan through 4 Feb (Bruce Smithson, m. obs.); at Glenburnie Quarry (restricted access), Craven Co, NC, 14 Feb (Ronnie Hewlette); and off North Topsail Beach, Onslow Co, NC, 28 Feb (Gilbert Grant).

Common Ground-Dove: One photographed along Baldwin Elkins Rd in Bladen Co, NC, 27 Jan (Amy Padgett) was the first sighting made in NC in three years.

White-winged Dove: One visiting a feeder in Wilmington, New Hanover Co, NC, 23 Dec through 23 Mar (Wanda King, m. obs.) was the only one reported this winter.



White-winged Dove 13 February 2018 New Hanover Co, NC Photo by Sam Cooper

Chuck-will's-widow: One was briefly seen roosting on the ground before being flushed in Sneads Ferry, Onslow Co, NC, after a locally unusual five-inch snowfall, 5 Jan (Gilbert Grant).

Nightjar sp.: One, probably either a lingering Chuck-will's-widow or a lingering Eastern Whip-poor-will, was seen in the campground at Pettigrew SP, Washington Co, NC, during its CBC, 27 Dec (Guy McGrane).

Ruby-throated Hummingbird: Several birders who hosted wintering Ruby-throated Hummingbirds in their yards at sites along the coast noted that some hummingbirds did not survive a period of persistent belowfreezing temperatures in early January (m. obs.).

Rufous Hummingbird: Sightings included an adult female visiting a feeder near Irmo, Lexington Co, SC, for a second winter in a row, 29 Nov into spring (Jerry Griggs); an adult male visiting a feeder in Black Mountain, Buncombe Co, NC, 4-19 Dec (Richard & Elizabeth Bruce); and an adult female visiting a feeder in Apex, Wake Co, NC, for a fourth winter in a row, banded 8 Feb (Susan Campbell).

Calliope Hummingbird: The immature male visiting a feeder in Charlotte, Mecklenburg Co, NC, 27 Nov into winter, was last seen 6 Jan (Jeff Lemons).

Broad-billed Hummingbird: The individual that visited a feeder in Southern Shores, Dare Co, NC, during the late fall, was found moribund after a night of below freezing temperatures, 1 Jan (Jeff Lewis).

Purple Gallinule: One photographed on Ocracoke Island, Hyde Co, NC, 15 Jan (*fide* Peter Vankevich) was unusual for the Carolinas in winter, especially for a site so far north.

Common Gallinule: 24 in Carteret Co, NC, during the Morehead City CBC, 17 Dec (*fide* John Fussell) was a good count for winter and the highest total on that CBC since 1984.

Sandhill Crane: Two seen in the Pungo Unit of Pocosin Lakes NWR, Washington Co, NC, from 12 Nov (Ricky Davis) were joined by a third crane 14 Dec (Greg Hudson) through 27 Feb (Stacy Barbour). Two wintered around the airport in Beaufort, Carteret Co, NC, again this year, 14 Dec through 8 Mar (Martin Wall, m. obs.). Two at Lake Junaluska, Haywood Co, NC, 1-2 Feb (m. obs.) were unusual for that site, probably forced down by inclement weather. A flock of nine was seen in flight over Cherokee Co, NC, near Farner, Tennessee, 17 Feb (Dennis Hammond). The high count of the wintering flock at the Bluff Unit of Santee NWR, Clarendon Co, SC, was 70 cranes on 21 Dec (Jay Chandler, et al.). 80 in flight over Lake Hartwell SP, Oconee Co, SC, 16 Feb (Brooks Garrett) was a notable count.



Sandhill Cranes 2 February 2018 Carteret Co, NC Photo by Martin Wall

Black-bellied Plover: One on the mudflats in the New Hope Creek arm of Jordan Lake, Chatham Co, NC, 3 Dec (Matt Spangler) and 4 Dec (Mark Kosiewski) was late for a site so far inland.

Long-billed Curlew: Two were seen around Raccoon Key, Cape Romain NWR, Charleston Co, SC, 17 Dec (John Cox, et al.) and 25 Feb (Caroline Eastman, et al.).

Hudsonian Godwit: One seen and photographed amongst Marbled Godwits at North Pond, Pea Island NWR, Dare Co, NC, 14 Dec (Peggy Eubank, Audrey Whitlock) was quite late.

Ruff: A female (Reeve) was seen and photographed in a marsh along US-17 near the Savannah Spoil Site, Jasper Co, SC, 19-24 Feb (Roger Smith, m. obs.).

Long-billed Dowitcher: 825 at the Savannah Spoil Site (restricted access), Jasper Co, SC, 30 Dec (Steve Calver) was a notable count.

American Woodcock: Several birders noted concentrations of woodcocks at coastal sites during a period of abnormal cold and snow in early January. Those sightings included 46+ around Oriental, Pamlico Co, NC, 6 Jan (Greg Perry); 22 video-recorded in a yard in Point Harbor, Currituck Co, NC, 6 Jan (*fide* Jeff Lewis); 15 at North River Preserve (limited access), Carteret Co, NC, 7 Jan (John Fussell, et al.); and 10 along the entrance road to the Bodie Island Lighthouse, Dare Co, NC, 7 Jan (Lewis).

Solitary Sandpiper: One photographed in Georgetown, Georgetown Co, SC, 12 Dec (Ritch Lilly) was late.

Red Phalarope: Sightings made from shore included one off Cape Point, Dare Co, NC, 14 Jan (Bruce Young, et al.); one photographed off Huntington Beach SP, Georgetown Co, SC, 16 Jan (Ritch Lilly); and two off Ft Fisher, New Hanover Co, NC, 27 Jan (Ricky Davis).

Great Skua: Remarkably, this species was seen on all seven pelagic trips in Dare Co, NC, this winter, with one 29 Dec, three 27 Jan, two 3 Feb, two 10 Feb, one 18 Feb, one 19 Feb, and two 24 Feb (Brian Patteson, et al.).

Pomarine Jaeger: One seen and photographed from the ocean overlook at Ft Fisher, New Hanover Co, NC, 2 Dec (Jamie Adams, Sam Cooper) was a good sighting from shore.

Parasitic Jaeger: Six in Carteret Co, NC, during the Morehead City CBC, 17 Dec (Mike Tove, et al.) was the highest total ever made on that count. Six off the Coquina rock outcrop at Ft Fisher, New Hanover Co, NC, 2 Dec (Jamie Adams, Sam Cooper) was a good total made from shore.

Dovekie: One around Johnnie Mercer's Pier, New Hanover Co, NC, 8 Feb (Jamie Adams) and 14 Feb (Sam Cooper) was a good find for a site south of the Outer Banks. 25 on a pelagic trip out of Hatteras, Dare Co, NC, 3 Feb (Brian Patteson, et al.) was the high count during the alcid irruption this winter.

Common Murre: 13 on a pelagic trip out of Wanchese, Dare Co, NC, 10 Feb (Brian Patteson, et al.) provided a new record high count of the species in NC. Common Murres were also seen on two pelagic trips out of Hatteras, albeit in smaller numbers, with four on 18 Feb and three on 19 Feb (Patteson, et al.)

Thick-billed Murre: Three were seen from shore this winter individuals off Coquina Beach, Dare Co, NC, 21 Jan (Ricky Davis); at Masonboro Inlet, New Hanover Co, NC, 21-22 Jan (Sherry Lane, m. obs.); and around the Kure Beach Pier, New Hanover Co, 11 Feb (Robert Snowden). On pelagic trips in Dare Co, NC, one was seen 10 Feb, two were seen 18 Feb, and one was seen 19 Feb (Brian Patteson, et al.).

Razorbill: There was a large movement of alcids off the NC coast in late January through mid-February, including a notable count of 12,000 Razorbills in flight off Cape Point, Dare Co, NC, 30 Jan (Brian Patteson).

Atlantic Puffin: This alcid was seen on all six pelagic trips in Dare Co, NC, in late January through February, with 15 on 27 Jan, five on 3 Feb, 17 on 10 Feb, three on 18 Feb, four on 19 Feb, and one on 24 Feb (Brian Patteson, et al.).

Black-legged Kittiwake: Sightings made on pelagic trips in Dare Co, NC, were five immature birds on 27 Jan, one immature bird on 3 Feb, five immature birds on 10 Feb, and one adult on 18 Feb (Brian Patteson, et al.). An immature bird in Wrightsville Beach, New Hanover Co, NC, 17 Feb (Jamie Adams, m. obs.) was the only one reported from shore this winter.



Black-legged Kittiwake 17 February 2018 New Hanover Co, NC Photo by Sam Cooper

Black-headed Gull: An adult at Mason Inlet, New Hanover Co, NC, 8 Jan intermittently through 27 Feb (Jamie Adams, m. obs.) was the only one reported this winter.

Little Gull: This species put on a good showing on pelagic trips in Dare Co, NC, in February, with two on 3 Feb, four on 10 Feb, one on 18 Feb, and three on 24 Feb (Brian Patteson, et al.). An adult photographed at Cape Point, Dare Co, NC, 17 Feb (Patteson) and an immature bird photographed in Wrightsville Beach, New Hanover Co, NC, 18 Feb (Jelmer Poelstra, m. obs.) were the only ones reported from shore.

Laughing Gull: An adult at Lake Johnson, Wake Co, NC, 25 Dec (Harry LeGrand) was unusual for a site in the Piedmont during the winter.

Ring-billed Gull: Unusual for the northern mountains in winter were four in West Jefferson, Ashe Co, NC, 8 Dec (Guy McGrane) and six in Boone, Watauga Co, NC, 8 Dec (Steve Dowlan).

Iceland Gull: This gull was well-reported again this winter. Sightings included two first-winter birds at Cape Point, Dare Co, NC, 3-10 Dec (Nathan Gatto, m. obs.); one first-winter bird at Cape Point 13 Jan into March (Jamie Adams, m. obs.); one first-winter bird at Murrells Inlet, Georgetown Co, SC, 28 Dec through 21 Jan (Sam Jolly, m. obs.); two on a pelagic trip out of Hatteras, Dare Co, NC, 29 Dec (Brian Patteson, et al.); one second-winter bird at Harris Lake, Wake Co, NC, 3 Jan (Lucas Bobay); an adult photographed on Lake Norman in Mecklenburg Co, NC, 6 Jan (Jeff Lemons, et al.); an adult on a pelagic trip out of Hatteras, 27 Jan (Brian Patteson, et al.); and a first-winter bird on Waites Island, Horry Co, SC, 6 Feb (Chris Hill, et al.). Lesser Black-backed Gull: 112 at Buckhorn Reservoir, Wilson Co, NC, 9 Feb (Ricky Davis) and 39 at Jordan Lake, Chatham Co, NC, 25 Feb (Jelmer Poelstra) were notable counts for those sites away from the coast. Individuals photographed on Lake Norman in Mecklenburg Co, NC, 7 Jan (Jeff Lemons, et al.) and along Milliken Rd in Spartanburg Co, SC, 22 Jan (Gary Harbour, Michael Robertson) were great finds for those sites in the western half of the Carolinas.

Glaucous Gull: Sightings included an immature bird in Williston, Carteret Co, NC, 18 Jan (Kathy Calvert); a first-winter bird at the Horry County Landfill (limited access), Horry Co, SC, 23-24 Jan (Chris Hill, m. obs.); an immature bird at Cape Point, Dare Co, NC, 24 Jan (Brian Patteson) and 25 Feb (John Haire); an adult photographed at Cape Point, 28 Jan (Lucas Bobay, Ed Corey); and a second-winter bird on a pelagic trip out of Hatteras, Dare Co, NC, 24 Feb (Brian Patteson, et al.).

Great Black-backed Gull: Farthest inland this winter was an adult at Buckhorn Reservoir, Wilson Co, NC, 9 Feb (Ricky Davis).

Red-billed Tropicbird: One of the most surprising sightings this winter was that of an adult Red-billed Tropicbird photographed flying south off the end of Jennette's Pier, Dare Co, NC, 9 Feb (David & Susan Disher, Marbry Hopkins). This sighting was the first ever made of this species in the Carolinas during the winter period.

Red-throated Loon: Inland, individuals were seen on Lake Hartwell, Anderson Co, SC, 14-19 Dec (Scott Davis, m. obs.); on Lake Norman in Mecklenburg Co, NC, 7 Jan (Jeff Lemons, et al.); and on Jordan Lake, Chatham Co, NC, 3 Feb (David Anderson, et al.).

Pacific Loon: Only one was reported this winter—an individual around Johnnie Mercer's Pier, New Hanover Co, NC, 22-27 Jan (Jamie Adams, m. obs.).

Northern Fulmar: This species was seen on all six pelagic trips in Dare Co, NC, in late January and in February, with a high count of 36 on 24 Feb (Brian Patteson, et al.).

Black-capped Petrel: Rarely seen in winter, two were seen on a pelagic trip out of Wanchese, Dare Co, NC, 10 Feb, and one was seen on a pelagic trip out of Hatteras, Dare Co, NC, 18 Feb (Brian Patteson, et al.)

Sooty Shearwater: Unusual for winter was one photographed over the ocean from Cape Point, Dare Co, NC, 2 Feb (Nathan Gatto, et al.) and two seen on a pelagic trip out of Wanchese, Dare Co, NC, 10 Feb (Brian Patteson, et al.).

Great Shearwater: Pelagic trips out of Hatters, Dare Co, NC, found one 27 Jan, two 3 Feb, and one 18 Feb (Brian Patteson, et al.).

Great Cormorant: Sightings included up to five at Masonboro Inlet, New Hanover Co, NC, throughout the winter (m. obs.); three at New River Inlet, Onslow Co, NC, 14 Dec (Gilbert Grant); and up to two at Beaufort Inlet, Carteret Co, NC, 17 Dec into spring (m. obs.)

Anhinga: Three lingered in Greenville, Pitt Co, NC, until its CBC, 2 Jan (*fide* Veronica Pantelidis), north of the species' typical winter range.

American White Pelican: A few of the higher counts this winter were 242 at Donnelley WMA, Colleton Co, SC, 14 Dec (David Perry); 227 at the Savannah Spoil Site (restricted access), Jasper Co, SC, 14 Dec (Steve Calver); 83 at Huntington Beach SP, Georgetown Co, SC, 21 Dec (Ritch Lilly); 120 at Bear Island WMA, Colleton Co, SC, 8 Jan (Patrick Markham); and a total of 75 at two sites near Davis, Carteret Co, NC, 20 Jan (Martin Wall). Six in Barnwell Co, SC, during the Savannah River Site CBC, 14 Dec (Peter Stangel) were new to that count.

Brown Pelican: One photographed at Robinwood Lake in Gaston Co, NC, during the Gastonia CBC, 16 Dec (Christy Hill, David Schroder, Steve Tracy) was a great find for a site so far from the coast.

Least Bittern: One seen and photographed at Savannah NWR, Jasper Co, SC, 22 Dec through 14 Jan (David Youngblood, m. obs.) was unusual for winter.

Reddish Egret: One seen along the Pitt St Causeway in Charleston Co, SC, 11-24 Feb (Takayuki Uchida, m. obs.) was unusual for winter.

Green Heron: One in Kitty Hawk, Dare Co, NC, during it CBC, 16 Dec (Lee Adams, Peggy Eubank) was unusual for a site so far north in winter.

Black-crowned Night-Heron: One at the Goldsboro WTP, Wayne Co, NC, during its CBC, 23 Dec (Mae Howell, David Williams) was locally unusual.

Roseate Spoonbill: 27 in Georgetown Co, SC, during the Winyah Bay CBC, 14 Dec (*fide* Lex Glover) was a notable count for winter and record number for that count.

Swallow-tailed Kite: One photographed over South Pond, Pea Island NWR, 24 Feb (Morgan Wilson) was quite early.

Swainson's Hawk: This species was seen at Alligator River NWR, Dare Co, NC, again this winter, with a light-morph immature bird seen 30 Dec (Audrey Whitlock, et al.) through 5 Feb (Jim Gould) and a light-morph adult bird seen 19 Jan (Guy McGrane, et al.) through 4 Feb (Jeff Lemons).

Red-tailed Hawk: A leucistic individual was seen by many at Alligator River NWR, Dare Co, NC, 16 Dec (Cindy Hamilton) through 10 Mar (Kevin Hudson).

Rough-legged Hawk: A light-morph immature bird was seen and photographed by many at Alligator River NWR, Dare Co, NC, 2 Jan (Jamie Adams) through 3 Mar (Alan MacEachren).

Golden Eagle: Sightings included an immature bird at Pee Dee NWR, Anson Co, NC, during its CBC, 30 Dec (Jennifer Daggy, et al.); an immature bird at Huntington Beach SP, Georgetown Co, SC, 13 Jan (Frank Lawkins) and 2 Feb (Ritch Lilly); a second-year bird photographed along Cheek Mountain Rd in Alleghany Co, NC, 21 Jan (Jesse Anderson, et al.); an adult over Seabrook Island, Charleston Co, SC, 15 Jan and 10 Feb (David Gardner, et al.); and an adult seen and photographed at Alligator River NWR, Dare Co, NC, 11-17 Feb (Rich DeMidio, m. obs.).

Snowy Owl: A minor irruption occurred this winter, with seven sightings made. One found at Pee Dee NWR, Anson Co, NC, 4 Dec, was taken to a wildlife rehabilitator but later died (*fide* Carolina Raptor Center). Individuals found in Myrtle Beach, Horry Co, SC, 12 Dec (*fide* Ritch Lilly) and at Piedmont Triad International Airport, Guilford Co, NC, in mid-December (*fide* Elizabeth Link) were collected by wildlife rehabilitators, treated and later released. One was seen around Pea Island NWR, Dare Co, NC, 15-17 Dec (Nathan & Sarah Gatto, m. obs.), 9 Jan (Becky Harrison, et al.), and 9 Feb (Karen Lebing). Because of the length of time between those sightings, it's likely the sightings represented three different individuals. One photographed at Lake Murray, Lexington Co, SC, 6 Jan (Julian Wilson) was not seen thereafter. All sightings made in SC are pending review by the SC BRC.

Short-eared Owl: One seen along Riverside Circle in Rockingham Co, NC, 1 Jan (Martin Wall) was a first for that county. Four on Marsh Island, Cape Romain NWR, Charleston Co, SC, 3 Feb (Jack Rogers) was a notable count.

Red-cockaded Woodpecker: 3+ were heard at North River Game Land in Currituck Co, NC, 16 Jan (Nick Flanders), near the northeast limit of the species' range.

Vermilion Flycatcher: Two were seen and photographed in SC this winter—a female in Georgetown, Georgetown Co, 23 Nov through 2 Jan (Ritch Lilly, m. obs.) and a male at Bookhart Farm in Orangeburg Co, 6 Dec through 2 Jan (Harold Donnelly, m. obs.).

Ash-throated Flycatcher: Individuals were seen on the west side of Lake Phelps, Washington Co, NC, during the Pettigrew SP CBC, 27 Dec (Al Hooks, Guy McGrane) and at the Savannah Spoil Site (restricted access), Jasper Co, SC, 30 Dec (Steve Calver, et al.).

Western Kingbird: Individuals were seen at North River Preserve (limited access), Carteret Co, NC, 3 Dec (Carol Reigle, et al.) and 31 Dec (Chandra Biggerstaff, et al.); in Oak Island, Brunswick Co, NC, during the Southport-Bald Head-Oak Islands CBC, 31 Dec (Matt Janson, Taylor Piephoff); and on Seabrook Island, Charleston Co, SC, 2-10 Jan (David Gardner, m. obs.).

Western Kingbird 3 December 2017 Carteret Co, NC Photo by Martin Wall



White-eyed Vireo: Six along Little Deep Creek Rd in the Croatan NF, Carteret Co, NC, 11 Dec (John Fussell), with four remaining at that site 23 Feb (Fussell), was a good count for that area in winter. One photographed at the Walnut Creek Wetland Center, Wake Co, NC, 16 Dec

(Edward Landi, et al.) was unusual for the Piedmont in winter.



White-eyed Vireo 7 February 2018 Carteret Co, NC Photo by Martin Wall

Warbling Vireo: One seen along Pine Grove Lane in Point Harbor, Currituck Co, NC, 13 Dec (Steve Ritt) was incredibly late. There is only one accepted winter report of this species in the Carolinas, one on Roanoke Island during the winter of 2001-2002.

Northern Rough-winged Swallow: One at Lake Crabtree, Wake Co, NC, 10 Dec (Harry LeGrand, Eddie Owens) was quite late.

Gray Catbird: One at Brookshire Park, Watauga Co, NC, 2 Dec (Guy McGrane, et al.) was late for a site in the mountains.

Snow Bunting: Sightings continued around the jetty on the south side of Oregon Inlet, Dare Co, NC, from 18 Nov to 17 Dec, with five seen 10 Dec (Jeff Beane, et al.).

Bachman's Sparrow: Counts of five at two sites in Croatan NF in Carteret Co, NC—along Millis Rd, 4 Feb (Martin Wall, et al.) and along Pringle Rd, 10 Feb (John Fussell)—were good for winter. Fussell notes that both sites had controlled burns the previous year.

Clay-colored Sparrow: Sightings, interestingly all made in Carteret Co, NC, this winter, were of two at North River Preserve (limited access), 3 Dec (John Fussell, et al.); one at North River Preserve 10 Dec (Martin Wall, et al.) and 31 Dec (Fussell, et al.); one at North River Golf Club, 14 Dec through 29 Jan (Fussell, m. obs.); and two at North River Golf Club, 29 Jan (Wall).

Vesper Sparrow: 15 at Donnelley WMA, Colleton Co, SC, 15 Feb (David Gardner, et al.) was a good count.

Lark Sparrow: Two were found at Archie Elledge WTP, Forsyth Co, NC, during the Winston-Salem CBC, 16 Dec (Nathan Gatto, et al.), with one continuing at that site through 15 Mar (m. obs.). One visited a feeder in Camden Co, NC, where photographed, 22 Feb into spring (James Harrison).

Henslow's Sparrow: One photographed in Croatan NF in Craven Co, NC, 5 Jan (Andrew Sodergren) was a good find for NC in winter.

LeConte's Sparrow: Two near the visitor center at Mackay Island NWR, Currituck Co, NC, 22 Dec (Eric Alton, Tamara Conklin) were unusual for the northeast corner of the Carolinas.

Lincoln's Sparrow: About a dozen were found across the Carolinas this winter. Three in a large industrial clear-cut with grass cover in Carteret Co, NC, 10 Dec (John Fussell, et al.) was a notable concentration.

Yellow-breasted Chat: Eight were reported in the Carolinas this winter, farthest inland were individuals visiting feeders in Mecklenburg Co, NC, 13-23 Dec (Taylor Piephoff) and in Spartanburg Co, SC, 30 Dec through 17 Feb (Susan Illner).

Yellow-headed Blackbird: Individuals were seen at a feeder in Pitt Co, NC, 4 Jan (Susan Bowen Moore); at a feeder in Chocowinity, Beaufort Co, NC, where photographed, 5 Jan (Ami Flowers Staples); in a yard in Morehead City, Carteret Co, NC, 9 Jan (Becky Brown); in Hobucken, Pamlico Co, NC, 10 Jan (Sheryl McNair, Christine Stoughton-Root); and along Baldwin Elkins Rd in Bladen Co, NC, where photographed, 26 Jan (Amy Padgett).

Bobolink: One photographed at Kinloch Plantation (restricted access), Georgetown Co, SC, during the Winyah Bay CBC, 14 Dec (Pam Ford, Craig Watson) was a truly amazing find, as there's only one previous winter record of the species in the Carolinas.

Bullock's Oriole: Two were seen in the Carolinas this winter—a first-year male visiting a feeder in Charlotte, Mecklenburg Co, NC, 19 Jan through 31 Mar (Carole Outwater, m. obs.) and an adult female visiting a feeder in Mt Pleasant, Charleston Co, SC, 22 Jan through 29 Mar (Dennis Forsythe, m. obs.), banded 9 Feb (Aaron Given).

Brewer's Blackbird: For a third winter in a row, this species was found in a horse pasture along Gum Swamp Rd in Hyde Co, NC, 29 Dec (John Fussell) through 20 Feb (Ron Clark), with a high count of ten at the site, 8 Feb (Jeff Lewis).

Tennessee Warbler: One seen and photographed along the Pitt St Causeway in Charleston Co, SC, 26-28 Jan (Elizabeth Anderegg, m. obs.) was noteworthy, as there are only five previous winter records for the Carolinas.

Nashville Warbler: One was seen in Oak Island, Brunswick Co, NC, during the Southport- Bald Head- Oak Islands CBC, 31 Dec (Matt Janson, Taylor Piephoff) and 1 Jan, when photographed (Jamie Adams).

American Redstart: Lingering into winter were individuals at North River Preserve (limited access), Carteret Co, NC, 10 Dec (John Fussell, et al.); in Gaston Co, NC, during the Gastonia CBC, 16 Dec (Ron Clark, Tom Sanders); and at Samworth WMA, Georgetown Co, SC, during the Litchfield-Pawleys Island CBC, 29 Dec (Irvin Pitts, Shawn Smolen-Morton)

Cape May Warbler: A male individual visited a suet feeder in Salisbury, Rowan Co, NC, where photographed, 24 Jan through 20 Feb (Steve Coggin).

Northern Parula: Lingering individuals were seen in Georgetown Co, SC, during the Winyah Bay CBC, 14 Dec (*fide* Lex Glover) and on Roanoke Island, Dare Co, NC, during the Kitty Hawk CBC, 16 Dec (Jeff Lewis).

Magnolia Warbler: One photographed in a yard in Charleston Co, SC, 8 Jan (Arch McCallum) was a great find as there are only three previous winter records in the Carolinas.

Black-throated Blue Warbler: A male individual in Myrtle Beach, Horry Co, SC, 2 Dec (Dick & Dorothy Rosche) was late to depart.

Yellow-rumped Warbler: 200 in central Halifax Co, NC, 28 Feb (Frank Enders) was a notable count for a site away from the coast.

Yellow-throated Warbler: Farthest inland this winter were individuals visiting feeders in Chapel Hill, Orange Co, NC, mid-December through 24 Jan (Tom Driscoll) and near Irmo, Lexington Co, SC, 5-10 Jan (Jerry Griggs).

Prairie Warbler: About a dozen were seen in the Carolinas this winter, all at sites along the coast. The northernmost was one at Mackay Island NWR, Currituck Co, NC, 10 Dec (Eric Alton, m. obs.) and two on Ocracoke Island, Hyde Co, NC, during its CBC, 30 Dec (*fide* Peter Vankevich).

Wilson's Warbler: Five were reported in the Carolinas this winter—an adult male in Point Harbor, Currituck Co, NC, 13 Dec (Steve Ritt); a female in a yard in Mt Pleasant, Charleston Co, SC, 13 Dec (John Cox); a male in Mt Pleasant, 28 Dec (Cox); a moribund individual in Morehead City, Carteret Co, NC, 30 Dec (Dave Lovett); and one photographed in Wilmington, New Hanover Co, NC, 7 Jan (Jamie Adams).

Summer Tanager: A female wintered around a yard in Brunswick Co, NC, for the fourth year in a row, 28 Dec into spring (Greg Massey). A female was seen near Merritt, Pamlico Co, NC, 15 Dec (Christine Stoughton-Root). An adult male was photographed at a feeder in Beaufort Co, SC, during the Hilton Head Island CBC, 15 Dec (*fide* Susan Murphy).

Western Tanager: This vagrant was very well-reported this winter. Sightings included up to two, an adult male and a bright-morph female, visiting a feeder in Charleston Co, SC, 30 Dec into March (Lacy & Russell Johnson, m. obs.); an adult male visiting a feeder on James Island, Charleston Co, SC, 3-30 Jan (Jennifer Tyrell, m. obs.); an adult male at Old Santee Canal Park, Berkley Co, SC, 5-17 Jan (Brad Sale, m. obs.); an adult male visiting a feeder in Scotts Hill, Pender Co, NC, 6-20 Jan (m. obs.); an adult male visiting a feeder in Wilmington, New Hanover Co, NC, 16 Jan (Bruce Smithson); a gray-morph female at Patriot's Point, Charleston Co, SC, 16 Jan (John Cox); a gray-morph female visiting a feeder on James Island, Charleston Co, SC, 11 Feb (Shannon Howard); and an adult male visiting a feeder in Stedman, Cumberland Co, NC, 19-24 Feb (Ann Horne, m. obs.).

Rose-breasted Grosbeak: Very unusual for winter was an adult male seen and photographed at a feeder in Cedar Mountain, Transylvania

Co, NC, late December through 15 Jan (Gail Wilcox) and an immature male photographed at a feeder in Gloucester, Carteret Co, NC, 15 Jan (Barbara Pigott).

Indigo Bunting: A male visiting a millet feeder in Meggett, Charleston Co, SC, where photographed, 13-16 Feb (Cherrie Sneed) must have been over-wintering in the area, as it was too early for a returning spring migrant.

Painted Bunting: 10+ in a yard in Straits, Carteret Co, NC, in February (Carolyn & Don Hoss) was a good count for a site so far north in winter. An adult male photographed at a feeder in Greenville, Pitt Co, NC, 6 Jan (Jerry Lotterhos) was locally unusual.

Dickcissel: A first-winter female was seen and photographed visiting a feeder at Myrtle Beach SP, Horry Co, SC, 30 Dec into spring (Scott Hartley, m. obs.).

Corrigenda: The Anna's Hummingbird that was banded in Buxton, Dare Co, NC, 16 Jan 2017, was first discovered in that yard the previous day, 15 Jan 2017, not 21 Dec 2016 as stated in the Winter 2017-18 "Briefs for the Files."



CAROLINA BIRD CLUB

www.carolinabirdclub.org

The Carolina Bird Club is a non-profit organization which represents and supports the birding community in the Carolinas through its official website, publications, meetings, workshops, trips, and partnerships, whose mission is

- To promote the observation, enjoyment, and study of birds.
- To provide opportunities for birders to become acquainted, and to share information and experience.
- To maintain well-documented records of birds in the Carolinas.
- To support the protection and conservation of birds and their habitats and foster an appreciation and respect of natural resources.
- To promote educational opportunities in bird and nature study.
- To support research on birds of the Carolinas and their habitats.

Membership is open to all persons interested in the conservation, natural history, and study of wildlife with particular emphasis on birds. Dues, contributions, and bequests to the Club may be deductible from state and federal income and estate taxes to the extent allowable. Pay dues or make donations at https://carolinabirdclub.org/dues. Make change of address at https://www.carolinabirdclub.org/members/profile. Send correspondence regarding membership matters to the Headquarters Secretary. Answers to questions about the club might be found at https://carolinabirdclub.org/about.html.

ANNUAL DUES

| Individual or non-profit | \$30.00 |
|--|----------|
| Family | \$35.00 |
| Student | |
| Sustaining and businesses | |
| Patron | |
| Life Membership (payable in four consecutive \$100 installments) | \$500.00 |
| Associate Life Membership (in same household as life member) | |

ELECTED OFFICERS

| President NC Vice-Presidents | Christine Stoughton-Root, Merritt NC Steve Tracy, Gastonia NC Sherry Lane, Wilmington NC | cssrbdr@gmail.com | | | |
|---------------------------------|--|-------------------|--|--|--|
| SC Vice-President | Craig Watson, Mt. Pleasant SC | | | | |
| Secretary | Jill Midgett, Charleston SC | | | | |
| Treasurer | Paul Dayer, Durham NC | | | | |
| NC Members-at-Large | Lester Coble, Ernul NC | | | | |
| | Karyl Gabriel, High Point NC | | | | |
| | Katherine Higgins, Wilmington NC | | | | |
| | Marilyn Westphal, Hendersonville NC | | | | |
| SC Members-at-Large | Mac Williams, Hartsville SC | | | | |
| | Steve McInnis, Columbia SC | | | | |
| | | | | | |

EX-OFFICIO EXECUTIVE COMMITTEE MEMBERS

Chat EditorSteve Shultz, Apex, NCchat@carolinabirdclub.orgNewsletter EditorVivian Glover, Orangeburg, SC newsletter@carolinabirdclub.orgWebsite EditorKent Fiala, Hillsborough, NCwebeditor@carolinabirdclub.orgImmediate Past PresidentIrvin Pitts, Lexington, NC

HEADQUARTERS SECRETARY

Carol Bowman

9 Quincy Place Pinehurst NC 28374 hq@carolinabirdclub.org